Prevent the spread of COVID-19 if you are sick

https://www.vdh.virginia.gov/epidemiology/respiratory%20diseases%20in%20virginia/precautions%20when%20sick/

If you are sick with COVID-19 or think you might have COVID-19, follow the steps below to care for yourself and to help protect other people in your home and community.

Stay home except to get medical care.

 Stay home. Most people with COVID-19 have mild illness and are able to recover at home without medical care. Do not leave your home, except to get medical care. Do not visit public areas.



- Take care of yourself. Get rest and stay hydrated. Take overthe-counter medicines, such as acetaminophen, to help you feel better.
- Stay in touch with your doctor. Call before you get medical care. Be sure to get care if you have trouble breathing, or have any other emergency warning signs, or if you think it is an emergency.
- Avoid public transportation, ride-sharing, or taxis.

Separate yourself from other people and pets in your home.

- As much as possible, stay in a specific room and away from other people and pets in your home. Also, you should use a separate bathroom, if available. If you need to be around other people or animals in or outside of the home, wear a mask.
 - See COVID-19 and Animals if you have questions about pets: https://www.cdc.gov/coronavirus/2019-ncov/daily-life-coping/animals.html
 - Additional guidance is available for disinfecting your home (https://health.students.vcu.edu/media/student-affairssites/ushs/docs/DisinfectingYourHome.pdf) and large households (https://health.students.vcu.edu/media/studentaffairs-sites/ushs/docs/GuidanceforLargeHouseholds.pdf).

Monitor your symptoms.

 Symptoms of COVID-19 include fever, cough, and shortness of breath but other symptoms may be present as well.



 Follow care instructions from your healthcare provider and local health department. Your local health authorities will give instructions on checking your symptoms and reporting information.

When to Seek Emergency Medical Attention

Look for emergency warning signs* for COVID-19. If someone is showing any of these signs, seek emergency medical care immediately:

- Trouble breathing
- · Persistent pain or pressure in the chest
- New confusion
- Bluish lips or face
- · Inability to wake or stay awake


*This list is not all possible symptoms. Please call your medical provider for any other symptoms that are severe or concerning to you.

Call 911 or call ahead to your local emergency facility:
Notify the operator that you are seeking care for someone who
has or may have COVID-19.

Call ahead before visiting your doctor.

- Call ahead. Many medical visits for routine care are being postponed or done by phone or telemedicine.
- If you have a medical appointment that cannot be postponed, call your doctor's office, and tell them you have or may have COVID-19.

If you are sick, wear a mask over your nose and mouth.





- You don't need to wear the mask if you are alone. If you can't
 put on a mask (because of trouble breathing for example),
 cover your coughs and sneezes in some other way. Try to stay at
 least 6 feet away from other people. This will help protect the
 people around you.
- Masks should not be placed on young children under age 2 years, anyone who has trouble breathing, or anyone who is not able to remove the mask without help.

Cover your coughs and sneezes.

- Cover your mouth and nose with a tissue when you cough or sneeze.
- Throw used tissues in a lined trash can.
- Immediately wash your hands with soap and water for at least 20 seconds. If soap and water are not available, clean your hands with an alcohol-based hand sanitizer that contains at least 60% alcohol.

Clean your hands often.

- Wash your hands often with soap and water for at least 20 seconds. This is especially important after blowing your nose, coughing, or sneezing; going to the bathroom; and before eating or preparing food.
- Use hand sanitizer if soap and water are not available. Use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry.
- Soap and water are the best option, especially if your hands are visibly dirty.
- Avoid touching your eyes, nose, and mouth with unwashed hands.

Avoid sharing personal household items.

- Do not share dishes, drinking glasses, cups, eating utensils, towels, or bedding with other people in your home.
- Wash these items thoroughly after using them with soap and water or put them in the dishwasher.

Clean all "high-touch" surfaces everyday.

 Clean and disinfect high-touch surfaces in your "sick room" and bathroom. Let someone else clean and disinfect surfaces in common areas, but not your bedroom and bathroom.



 If a caregiver or other person needs to clean and disinfect a sick person's bedroom or bathroom, they should do so on an as-needed basis. The caregiver/other person should wear a mask and wait as long as possible after the sick person has used the bathroom.

High-touch surfaces include phones, remote controls, counters, tabletops, doorknobs, bathroom fixtures, toilets, keyboards, tablets, and bedside tables.

- Clean and disinfect areas that may have blood, stool, or body fluids on them.
- Use household cleaners and disinfectants. Clean the area or item with soap and water or another detergent if it is dirty. Then use a household disinfectant.
 - Be sure to follow the instructions on the label to ensure safe and effective use of the product. Many products recommend keeping the surface wet for several minutes to ensure germs are killed. Many also recommend precautions such as wearing gloves and making sure you have good ventilation during use of the product.
 - Most EPA-registered household disinfectants should be effective.

When You Can be Around Others After You Had or Likely Had COVID-19



When you can be around others (end home isolation) depends on different factors for different situations.

- I think or know I had COVID-19, and I had symptoms
 - You can be with others after
 - 24 hours with no fever
 AND
 - Symptoms improved

without use of a fever-reducing medication (ibuprofen/Advil/Motrin, acetaminophen/Tylenol, etc.).

 When returning to normal activities, wear a mask for the next 5 days when indoors and around others. You are less contagious during this time, but it is still important to be cautious and social distance, especially if you will be around those at higher risk for serious illness (people with diabetes, asthma, immune suppression; babies; pregnant individuals; elderly).